

**DIMENSIONAL REDUCTION, $SL(2, \mathbb{C})$ -EQUIVARIANT BUNDLES
AND STABLE HOLOMORPHIC CHAINS**

LUIS ÁLVAREZ-CÓNSUL

*Department of Mathematics, University of Illinois at Urbana-Champaign,
IL 61801, USA**E-mail: lalvarez@math.uiuc.edu*

OSCAR GARCÍA-PRADA

*Departamento de Matemáticas, Universidad Autónoma de Madrid,
28049 Madrid, Spain**E-mail: oscar.garcia-prada@uam.es*

In this paper we study gauge theory on $SL(2, \mathbb{C})$ -equivariant bundles over $X \times \mathbb{P}^1$, where X is a compact Kähler manifold, $\mathbb{P}^1$ is the complex projective line, and the action of $SL(2, \mathbb{C})$ is trivial on X and standard on $\mathbb{P}^1$. We first classify these bundles, showing that they are in correspondence with objects on X — that we call holomorphic chains — consisting of a finite number of holomorphic bundles E_i and morphisms $E_i \rightarrow E_{i-1}$. We then prove a Hitchin–Kobayashi correspondence relating the existence of solutions to certain natural gauge-theoretic equations and an appropriate notion of stability for an equivariant bundle and the corresponding chain. A central tool in this paper is a dimensional reduction procedure which allow us to go from $X \times \mathbb{P}^1$ to X .

Keywords: Equivariant bundle, stability, Hermitian-Einstein equations, vortex equations, holomorphic chains, Hitchin-Kobayashi correspondence, dimensional reduction.

Note: A complete PDF file can be obtained from <http://www.worldscinet.com/journals/ijm/>
See also <http://www.math.uiuc.edu/~lalvarez/papers/>